

FACTS *for* FAMILIES

No. 47

November 2012

The Anxious Child

All children experience anxiety. Anxiety in children is expected and normal at specific times in development. For example, from approximately age 8 months through the preschool years, healthy youngsters may show intense distress (anxiety) at times of separation from their parents or other persons with whom they are close. Young children may have short-lived fears, such as fear of the dark, storms, animals, or a fear of strangers. Anxious children are often overly tense or uptight. Some may seek a lot of reassurance, and their worries may interfere with activities. Parents should not dismiss a child's fears. Because anxious children may also be quiet, compliant and eager to please, their difficulties may be missed. Parents should be alert to the signs of severe anxiety so they can intervene early to prevent complications. There are different types of anxiety in children.

Symptoms of separation anxiety include:

- constant thoughts and intense fears about the safety of parents and caretakers
- refusing to go to school
- frequent stomachaches and other physical complaints
- extreme worries about sleeping away from home
- being overly clingy
- panic or tantrums at times of separation from parents
- trouble sleeping or nightmares

Symptoms of phobia include:

- extreme fear about a specific thing or situation (ex. dogs, insects, or needles)
- the fears cause significant distress and interfere with usual activities

Symptoms of social anxiety include:

- fears of meeting or talking to people
- avoidance of social situations
- few friends outside the family

Other symptoms of anxious children include:

- many worries about things before they happen
- constant worries or concerns about family, school, friends, or activities
- repetitive, unwanted thoughts (obsessions) or actions (compulsions)
- fears of embarrassment or making mistakes

The Anxious Child, “Facts for Families,” No. 47 (11/12)

- low self-esteem and lack of self-confidence

Severe anxiety problems in children can be treated. Early treatment can prevent future difficulties, such as loss of friendships, failure to reach social and academic potential, and feelings of low self-esteem. Treatments may include a combination of the following: individual psychotherapy, family therapy, medications, behavioral treatments, and consultation to the school.

If anxieties become severe and begin to interfere with the child’s usual activities (for example separating from parents, attending school and making friends) parents should consider seeking an evaluation from a qualified mental health professional or a child and adolescent psychiatrist.

If you find **Facts for Families**® helpful and would like to make good mental health a reality, consider donating to the [Campaign for America’s Kids](#). Your support will help us continue to produce and distribute Facts for Families, as well as other vital mental health information, free of charge.

You may also mail in your contribution. Please make checks payable to the AACAP and send to *Campaign for America’s Kids*, P.O. Box 96106, Washington, DC 20090.

The American Academy of Child and Adolescent Psychiatry (AACAP) represents over 8,500 child and adolescent psychiatrists who are physicians with at least five years of additional training beyond medical school in general (adult) and child and adolescent psychiatry.

Facts for Families© information sheets are developed, owned and distributed by AACAP. Hard copies of **Facts** sheets may be reproduced for personal or educational use without written permission, but cannot be included in material presented for sale or profit. All **Facts** can be viewed and printed from the AACAP website (www.aacap.org). **Facts** sheets may not be reproduced, duplicated or posted on any other website without written consent from AACAP. Organizations are permitted to create links to AACAP’s website and specific **Facts** sheets. For all questions please contact the AACAP Communications & Marketing Coordinator, ext. 154.

If you need immediate assistance, please dial 911.

Copyright © 2012 by the American Academy of Child and Adolescent Psychiatry.